

Fourier transforms of measures supported on graphs

Jonathan M. Fraser

joint work with **Tuomas Orponen** and **Tuomas Sahlsten**



My coauthors



My coauthors



Fourier analysis and Hausdorff dimension

Fourier analysis and Hausdorff dimension

Let $K \subset \mathbb{R}^2$ be Borel and $\mathcal{P}(K)$ be the set of all Borel probability measures on K .

Fourier analysis and Hausdorff dimension

Let $K \subset \mathbb{R}^2$ be Borel and $\mathcal{P}(K)$ be the set of all Borel probability measures on K .

$$\dim_{\text{H}} K = \sup \left\{ s \leq 2 : \exists \mu \in \mathcal{P}(K), I_s(\mu) := \iint \frac{d\mu(x)d\mu(y)}{|x-y|^s} < \infty \right\}$$

Fourier analysis and Hausdorff dimension

Let $K \subset \mathbb{R}^2$ be Borel and $\mathcal{P}(K)$ be the set of all Borel probability measures on K .

$$\dim_{\text{H}} K = \sup \left\{ s \leq 2 : \exists \mu \in \mathcal{P}(K), I_s(\mu) := \iint \frac{d\mu(x)d\mu(y)}{|x-y|^s} < \infty \right\}$$

- The **Fourier transform** of $\mu \in \mathcal{P}(K)$ is $\hat{\mu} : \mathbb{R}^2 \rightarrow \mathbb{C}$

$$\hat{\mu}(\xi) := \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} d\mu(x) \quad \xi \in \mathbb{R}^2$$

Fourier analysis and Hausdorff dimension

Let $K \subset \mathbb{R}^2$ be Borel and $\mathcal{P}(K)$ be the set of all Borel probability measures on K .

$$\dim_{\text{H}} K = \sup \left\{ s \leq 2 : \exists \mu \in \mathcal{P}(K), I_s(\mu) := \iint \frac{d\mu(x)d\mu(y)}{|x-y|^s} < \infty \right\}$$

- The **Fourier transform** of $\mu \in \mathcal{P}(K)$ is $\widehat{\mu} : \mathbb{R}^2 \rightarrow \mathbb{C}$

$$\widehat{\mu}(\xi) := \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} d\mu(x) \quad \xi \in \mathbb{R}^2$$

- Alternative formula for the s -energy:

$$I_s(\mu) = c \int_{\mathbb{R}^2} |\widehat{\mu}(\xi)|^2 |\xi|^{s-2} d\xi$$

Fourier analysis and Hausdorff dimension

Let $K \subset \mathbb{R}^2$ be Borel and $\mathcal{P}(K)$ be the set of all Borel probability measures on K .

$$\dim_{\text{H}} K = \sup \left\{ s \leq 2 : \exists \mu \in \mathcal{P}(K), I_s(\mu) := \iint \frac{d\mu(x)d\mu(y)}{|x-y|^s} < \infty \right\}$$

- The **Fourier transform** of $\mu \in \mathcal{P}(K)$ is $\hat{\mu} : \mathbb{R}^2 \rightarrow \mathbb{C}$

$$\hat{\mu}(\xi) := \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} d\mu(x) \quad \xi \in \mathbb{R}^2$$

- Alternative formula for the s -energy:

$$I_s(\mu) = c \int_{\mathbb{R}^2} |\hat{\mu}(\xi)|^2 |\xi|^{s-2} d\xi$$

...so if for $\varepsilon > 0$ we have

$$|\hat{\mu}(\xi)| \lesssim |\xi|^{-(s+\varepsilon)/2}, \xi \in \mathbb{R}^2$$

Fourier analysis and Hausdorff dimension

Let $K \subset \mathbb{R}^2$ be Borel and $\mathcal{P}(K)$ be the set of all Borel probability measures on K .

$$\dim_{\text{H}} K = \sup \left\{ s \leq 2 : \exists \mu \in \mathcal{P}(K), I_s(\mu) := \iint \frac{d\mu(x)d\mu(y)}{|x-y|^s} < \infty \right\}$$

- The **Fourier transform** of $\mu \in \mathcal{P}(K)$ is $\hat{\mu} : \mathbb{R}^2 \rightarrow \mathbb{C}$

$$\hat{\mu}(\xi) := \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} d\mu(x) \quad \xi \in \mathbb{R}^2$$

- Alternative formula for the s -energy:

$$I_s(\mu) = c \int_{\mathbb{R}^2} |\hat{\mu}(\xi)|^2 |\xi|^{s-2} d\xi$$

...so if for $\varepsilon > 0$ we have

$$|\hat{\mu}(\xi)| \lesssim |\xi|^{-(s+\varepsilon)/2}, \xi \in \mathbb{R}^2 \implies I_s(\mu) < \infty$$

Fourier analysis and Hausdorff dimension

Let $K \subset \mathbb{R}^2$ be Borel and $\mathcal{P}(K)$ be the set of all Borel probability measures on K .

$$\dim_{\text{H}} K = \sup \left\{ s \leq 2 : \exists \mu \in \mathcal{P}(K), I_s(\mu) := \iint \frac{d\mu(x)d\mu(y)}{|x-y|^s} < \infty \right\}$$

- The **Fourier transform** of $\mu \in \mathcal{P}(K)$ is $\hat{\mu} : \mathbb{R}^2 \rightarrow \mathbb{C}$

$$\hat{\mu}(\xi) := \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} d\mu(x) \quad \xi \in \mathbb{R}^2$$

- Alternative formula for the s -energy:

$$I_s(\mu) = c \int_{\mathbb{R}^2} |\hat{\mu}(\xi)|^2 |\xi|^{s-2} d\xi$$

...so if for $\varepsilon > 0$ we have

$$|\hat{\mu}(\xi)| \lesssim |\xi|^{-(s+\varepsilon)/2}, \xi \in \mathbb{R}^2 \implies I_s(\mu) < \infty \implies \dim_{\text{H}} K \geq s.$$

Fourier analysis and Hausdorff dimension

This motivates...

Definition (Fourier dimension)

$$\dim_{\mathbb{F}} K := \sup\{s \leq 2 : \exists \mu \in \mathcal{P}(K) \text{ with } |\widehat{\mu}(\xi)| \lesssim |\xi|^{-s/2}, \xi \in \mathbb{R}^2\}$$

Fourier analysis and Hausdorff dimension

This motivates...

Definition (Fourier dimension)

$$\dim_{\mathbb{F}} K := \sup\{s \leq 2 : \exists \mu \in \mathcal{P}(K) \text{ with } |\widehat{\mu}(\xi)| \lesssim |\xi|^{-s/2}, \xi \in \mathbb{R}^2\}$$

...and shows that $\dim_{\mathbb{F}} K \leq \dim_{\mathbb{H}} K$.

Fourier analysis and Hausdorff dimension

This motivates...

Definition (Fourier dimension)

$$\dim_{\mathbb{F}} K := \sup\{s \leq 2 : \exists \mu \in \mathcal{P}(K) \text{ with } |\widehat{\mu}(\xi)| \lesssim |\xi|^{-s/2}, \xi \in \mathbb{R}^2\}$$

...and shows that $\dim_{\mathbb{F}} K \leq \dim_{\mathbb{H}} K$.

Definition (Round sets)


If $\dim_{\mathbb{F}} K = \dim_{\mathbb{H}} K$, we say that K is **round**.

- Round sets are also known as **Salem sets**.

Finding round sets

Finding round sets


Unit circle S^1 is round...



...since $\widehat{\mathcal{H}^1 \llcorner S^1}$ decays like $|\xi|^{-1/2}$.


Finding round sets II

A line $L \subset \mathbb{R}^2$ is **not** round...



Finding round sets II

A line $L \subset \mathbb{R}^2$ is **not** round...




...since $\hat{\mu} \equiv 1$ on L^\perp for any $\mu \in \mathcal{P}(L)$!

$$\implies \dim_{\mathbb{F}} L = 0 < 1 = \dim_{\mathbb{H}} L.$$

Finding round sets II

A line $L \subset \mathbb{R}^2$ is **not** round...




...since $\hat{\mu} \equiv 1$ on L^\perp for any $\mu \in \mathcal{P}(L)$!

$$\implies \dim_{\mathbb{F}} L = 0 < 1 = \dim_{\mathbb{H}} L.$$

- Punchline: $\dim_{\mathbb{H}}$ measures size, but $\dim_{\mathbb{F}}$ also contains information on **curvature**.

Finding round sets II

A line $L \subset \mathbb{R}^2$ is **not** round...



...since $\hat{\mu} \equiv 1$ on L^\perp for any $\mu \in \mathcal{P}(L)$!


$$\implies \dim_{\mathbb{F}} L = 0 < 1 = \dim_{\mathbb{H}} L.$$

- Punchline: $\dim_{\mathbb{H}}$ measures size, but $\dim_{\mathbb{F}}$ also contains information on **curvature**.

'Non-trivial' round sets are hard to construct deterministically,

Finding round sets II

A line $L \subset \mathbb{R}^2$ is **not** round...



...since $\hat{\mu} \equiv 1$ on L^\perp for any $\mu \in \mathcal{P}(L)$!

$$\implies \dim_{\mathbb{F}} L = 0 < 1 = \dim_{\mathbb{H}} L.$$

- Punchline: $\dim_{\mathbb{H}}$ measures size, but $\dim_{\mathbb{F}}$ also contains information on **curvature**.

'Non-trivial' round sets are hard to construct deterministically, but

- there are some examples by Kahane and Kaufman; in particular some sets arising from Diophantine approximation are round.

Finding round sets III

There are many **random** round sets.

Finding round sets III

There are many **random** round sets. First such constructions are due to Salem, but the following key result is by Kahane.

Theorem (Kahane 1986)

Let $\omega: [0, \infty) \rightarrow \mathbb{R}$ be 1-dimensional Brownian motion, and let $K \subset [0, \infty)$ be compact. Then the image $\omega(K) \subset \mathbb{R}$ is a.s. round, with

$$\dim_{\mathbb{F}} \omega(K) = \dim_{\mathbb{H}} \omega(K) = \min\{1, 2 \dim_{\mathbb{H}} K\}.$$

Finding round sets III

There are many **random** round sets. First such constructions are due to Salem, but the following key result is by Kahane.

Theorem (Kahane 1986)

Let $\omega: [0, \infty) \rightarrow \mathbb{R}$ be 1-dimensional Brownian motion, and let $K \subset [0, \infty)$ be compact. Then the image $\omega(K) \subset \mathbb{R}$ is a.s. round, with

$$\dim_{\mathbb{F}} \omega(K) = \dim_{\mathbb{H}} \omega(K) = \min\{1, 2 \dim_{\mathbb{H}} K\}.$$

Analogous result also holds for fractional Brownian motion.

Finding round sets IV

So, the image of any compact set under a 'random function' is round.

Finding round sets IV

So, the image of any compact set under a 'random function' is round.

- Maybe random functions provide more examples of round sets?

Finding round sets IV

So, the image of any compact set under a 'random function' is round.

- Maybe random functions provide more examples of round sets?

Kahane writes (1993):

“...proving almost sure roundedness for specific random sets is never easy and it remains an open program for most natural random sets: level sets and graphs of random functions in particular.”

Finding round sets IV

So, the image of any compact set under a 'random function' is round.

- Maybe random functions provide more examples of round sets?

Kahane writes (1993):

"...proving almost sure roundedness for specific random sets is never easy and it remains an open program for most natural random sets: level sets and graphs of random functions in particular."

Later (2006), Shieh and Xiao explicitly ask:

"Are the graph and level sets of a stochastic process such as fractional Brownian motion Salem sets?"

Finding round sets V

The conjecture is partially confirmed for level sets:

Theorem (Fouché and Mukeru 2013)

Let $\omega : [0, \infty) \rightarrow \mathbb{R}$ be 1-dimensional fractional Brownian motion.
Then for $a \in \mathbb{R}$, the level set

$$L_\omega(a) = \{0 \leq t \leq 1 : \omega(t) = a\}$$

is round with positive probability and

$$\dim_{\text{F}} L_\omega(a) = \dim_{\text{H}} L_\omega(a) = \frac{1}{2}.$$

Graphs of Brownian motion

How about graphs of 1-dimensional (fractional) Brownian motion?

Graphs of Brownian motion

How about graphs of 1-dimensional (fractional) Brownian motion?

The Hausdorff dimension part is classical:

Theorem (Taylor 1953, Adler 1977)

Let $\omega : [0, \infty) \rightarrow \mathbb{R}$ be the 1-dimensional fractional Brownian motion with Hurst exponent $0 < H < 1$. Then, the graph

$$G_\omega := \{(t, \omega(t)) : t \in [0, \infty)\} \subset \mathbb{R}^2$$

a.s. satisfies

$$\dim_{\text{H}} G_\omega = 2 - H.$$

- In particular, $\dim_{\text{H}} G_\omega > 1$ a.s.

Fourier dimension of graphs

We proved:

Theorem (J. F., T. Orponen and T. Sahlsten. 2013)

Let $E \subset \mathbb{R}$ be a set, and let $f: E \rightarrow \mathbb{R}$ be a function. Then

$$\dim_{\mathbb{F}} G_f \leq 1.$$

Fourier dimension of graphs

We proved:

Theorem (J. F., T. Orponen and T. Sahlsten. 2013)

Let $E \subset \mathbb{R}$ be a set, and let $f: E \rightarrow \mathbb{R}$ be a function. Then

$$\dim_{\mathbb{F}} G_f \leq 1.$$

Combining this with Taylor's and Adler's results answers Kahane's, Shieh's and Xiao's questions on random graphs in the negative:

Corollary

The Brownian graphs G_ω are a.s. **not** round/Salem.

Proof

Proof

Some theorems in **geometric measure theory** can be strengthened if hypotheses on Hausdorff dimension are replaced by those on Fourier dimension.

Proof

Some theorems in **geometric measure theory** can be strengthened if hypotheses on Hausdorff dimension are replaced by those on Fourier dimension.

- **Marstrand's projection theorem:** if a planar set K has Fourier dimension $s \in [0, 1]$, then all projections of K onto lines have Fourier dimension $\geq s$ (folklore).

Proof

Some theorems in **geometric measure theory** can be strengthened if hypotheses on Hausdorff dimension are replaced by those on Fourier dimension.

- **Marstrand's projection theorem:** if a planar set K has Fourier dimension $s \in [0, 1]$, then all projections of K onto lines have Fourier dimension $\geq s$ (folklore).
- **Falconer's distance set conjecture:** if a planar set K has Fourier dimension $s > 1$, then the distance set of K , namely

$$\Delta(K) = \{|x - y| : x, y \in K\},$$

has positive length (P. Mattila).

Proof II

For us, the key is proving a Fourier analytic version of Marstrand's slicing theorem.

- **Marstrand's slicing theorem:** if a planar set K has $\dim_{\mathbb{F}} K > 1$, then in every direction there are Leb positively many lines ℓ with

$$\dim_{\mathbb{H}}[K \cap \ell] > 0.$$

Proof II

For us, the key is proving a Fourier analytic version of Marstrand's slicing theorem.

- **Marstrand's slicing theorem:** if a planar set K has $\dim_{\mathbb{F}} K > 1$, then in every direction there are Leb positively many lines ℓ with

$$\dim_{\mathbb{H}}[K \cap \ell] > 0.$$

In particular, the above conclusion holds for lines in the **vertical** direction.

Proof II


For us, the key is proving a Fourier analytic version of Marstrand's slicing theorem.

- **Marstrand's slicing theorem:** if a planar set K has $\dim_{\mathbb{F}} K > 1$, then in every direction there are Leb positively many lines ℓ with

$$\dim_{\mathbb{H}}[K \cap \ell] > 0.$$

In particular, the above conclusion holds for lines in the **vertical** direction.

- Graphs of arbitrary functions f clearly do not have this property:



Hence, they can have Fourier dimension *at most one!*

Proof III

- **To do:** inspect the proof of Marstrand's slicing theorem.

Proof III

- **To do:** inspect the proof of Marstrand's slicing theorem.
- **The enemy:** classical proofs do not use Fourier analysis (i.e. relation with Fourier transforms and dimension is not clear).

Proof III

- **To do:** inspect the proof of Marstrand's slicing theorem.
- **The enemy:** classical proofs do not use Fourier analysis (i.e. relation with Fourier transforms and dimension is not clear).
- **The solution:** invent a Fourier analytic proof for Marstrand's slicing theorem.

Proof IV

Proof sketch:

- Assume that $K \subset \mathbb{R}^2$ is a set with $\dim_{\mathbb{F}} K > 1$.

Proof IV



Proof sketch:

- Assume that $K \subset \mathbb{R}^2$ is a set with $\dim_{\mathbb{F}} K > 1$.
- Choose $\mu \in \mathcal{P}(K)$ with $|\widehat{\mu}(\xi)| \lesssim |\xi|^{-(s+\varepsilon)/2}$ for some $s > 1$.

Proof IV

Proof sketch:

- Assume that $K \subset \mathbb{R}^2$ is a set with $\dim_{\mathbb{F}} K > 1$.
- Choose $\mu \in \mathcal{P}(K)$ with $|\widehat{\mu}(\xi)| \lesssim |\xi|^{-(s+\varepsilon)/2}$ for some $s > 1$.
- **Slice** the measure μ with vertical lines $L_t = \{(t, y) : y \in \mathbb{R}\} \subset \mathbb{R}^2$ to obtain 'sliced measures' μ_t , supported on $K \cap L_t$.



Easy but important: $\mu_t \neq 0$ for Leb positively many t .
(this requires the decay assumption of $\widehat{\mu}$ and Plancherel's formula)

Proof V

- Consider the $(s - 1)$ -energies

$$I_{s-1}(\mu_t) = \iint \frac{d\mu_t(x) d\mu_t(y)}{|x - y|^{s-1}} \dots$$

Proof V

- Consider the $(s - 1)$ -energies

$$I_{s-1}(\mu_t) = \iint \frac{d\mu_t(x) d\mu_t(y)}{|x - y|^{s-1}} \dots$$

- ...and prove the inequality

$$\int_{\mathbb{R}} I_{s-1}(\mu_t) dt \lesssim \int_{\mathbb{R}^2} |\widehat{\mu}(\xi)|^2 |\xi_2|^{s-2} d\xi \dots$$

Proof V

- Consider the $(s - 1)$ -energies

$$I_{s-1}(\mu_t) = \iint \frac{d\mu_t(x) d\mu_t(y)}{|x - y|^{s-1}} \dots$$

- ...and prove the inequality

$$\int_{\mathbb{R}} I_{s-1}(\mu_t) dt \lesssim \int_{\mathbb{R}^2} |\widehat{\mu}(\xi)|^2 |\xi_2|^{s-2} d\xi \dots$$

- ...which follows from Plancherel if μ is a smooth function; the general case involves a tedious approximation.

Proof V

- Consider the $(s - 1)$ -energies

$$I_{s-1}(\mu_t) = \iint \frac{d\mu_t(x) d\mu_t(y)}{|x - y|^{s-1}} \dots$$

- ...and prove the inequality

$$\int_{\mathbb{R}} I_{s-1}(\mu_t) dt \lesssim \int_{\mathbb{R}^2} |\widehat{\mu}(\xi)|^2 |\xi_2|^{s-2} d\xi \dots$$

- ...which follows from Plancherel if μ is a smooth function; the general case involves a tedious approximation.
- In fact, this inequality was proved by Orponen (2012).

Proof V

- Consider the $(s - 1)$ -energies

$$I_{s-1}(\mu_t) = \iint \frac{d\mu_t(x) d\mu_t(y)}{|x - y|^{s-1}} \dots$$

- ...and prove the inequality

$$\int_{\mathbb{R}} I_{s-1}(\mu_t) dt \lesssim \int_{\mathbb{R}^2} |\widehat{\mu}(\xi)|^2 |\xi_2|^{s-2} d\xi \dots$$

- ...which follows from Plancherel if μ is a smooth function; the general case involves a tedious approximation.
- In fact, this inequality was proved by Orponen (2012).
- Plug in the estimate $|\widehat{\mu}(\xi)|^2 \lesssim |\xi|^{-(s+\varepsilon)}$ and check that the integral on the R.H.S is finite.

Proof V

- Consider the $(s - 1)$ -energies

$$I_{s-1}(\mu_t) = \iint \frac{d\mu_t(x) d\mu_t(y)}{|x - y|^{s-1}} \dots$$

- ...and prove the inequality

$$\int_{\mathbb{R}} I_{s-1}(\mu_t) dt \lesssim \int_{\mathbb{R}^2} |\widehat{\mu}(\xi)|^2 |\xi_2|^{s-2} d\xi \dots$$

- ...which follows from Plancherel if μ is a smooth function; the general case involves a tedious approximation.
- In fact, this inequality was proved by Orponen (2012).
- Plug in the estimate $|\widehat{\mu}(\xi)|^2 \lesssim |\xi|^{-(s+\varepsilon)}$ and check that the integral on the R.H.S is finite.

$$\implies I_{s-1}(\mu_t) < \infty \quad \text{for Lebesgue a.e. } t \in \mathbb{R}.$$

Proof V

- Consider the $(s - 1)$ -energies

$$I_{s-1}(\mu_t) = \iint \frac{d\mu_t(x) d\mu_t(y)}{|x - y|^{s-1}} \dots$$

- ...and prove the inequality

$$\int_{\mathbb{R}} I_{s-1}(\mu_t) dt \lesssim \int_{\mathbb{R}^2} |\widehat{\mu}(\xi)|^2 |\xi_2|^{s-2} d\xi \dots$$

- ...which follows from Plancherel if μ is a smooth function; the general case involves a tedious approximation.
- In fact, this inequality was proved by Orponen (2012).
- Plug in the estimate $|\widehat{\mu}(\xi)|^2 \lesssim |\xi|^{-(s+\varepsilon)}$ and check that the integral on the R.H.S is finite.

$$\implies I_{s-1}(\mu_t) < \infty \quad \text{for Leb a.e. } t \in \mathbb{R}.$$

$$\implies \dim_{\text{H}}[K \cap L_t] \geq s - 1 > 0 \quad \text{for Leb pos. many } t.$$

Q.E.D.

Open questions

- What is the a.s. Fourier dimension of the Brownian graphs G_ω ?
We only proved that $\dim_{\mathbb{F}} G_\omega \leq 1$.

Open questions

- What is the a.s. Fourier dimension of the Brownian graphs G_ω ?
We only proved that $\dim_{\mathbb{F}} G_\omega \leq 1$.
- Is Fourier dimension countably (or even finitely) stable? I.e.

$$\dim_{\mathbb{F}} \left(\bigcup_{i \in \mathbb{N}} A_i \right) = \sup_i \dim_{\mathbb{F}} A_i?$$

Further results

Not being able to solve the first open question, we considered the following variant:

Question

What is the Fourier dimension of the graph of a typical function $f \in C[0, 1]$?

- Here we mean *typical* in the sense of Baire category.

Further results II

We proved:

Theorem (J. F., T. Orponen and T. Sahlsten. 2013)

The typical function $f \in C[0, 1]$ has the following property. If $\mu \in \mathcal{P}(G_f)$, then

$$\limsup_{|\xi| \rightarrow \infty} |\widehat{\mu}(\xi)| \geq \frac{1}{5}.$$

In particular, $\dim_{\mathbb{F}} G_f = 0$.

- The constant $1/5$ is not sharp (optimal constant unknown).
- Hausdorff dimension $\dim_{\mathbb{H}} G_f \geq 1$ for any $f \in C[0, 1]$, so our result implies that the graph of a typical function is *not* round/Salem!

References

J. M. Fraser, T. Orponen, T. Sahlsten: On Fourier analytic properties of graphs, to appear in *Int. Math. Res. Not. IMRN*, (2013), (arXiv:1211.4803v2).

W. Fouch, S. Mukeru: On the Fourier structure of the zero set of fractional Brownian motion, *Statistics & Probability Letters* **83**:2 (2013), 459–466.

J.-P. Kahane: Fractals and random measures, *Bull. Sci. Math.*, **117** (1993), 153–159.

T. Orponen: Slicing sets and measures, and the dimension of exceptional parameters, *J. Geom. Anal.*, **24**, (2014), 47–80.

N.-R. Shieh, Y. Xiao: Images of Gaussian random fields: Salem sets and interior points, *Studia Math.*, **176**, (2006), 37–60.